

Specification

Liquid medium for the selective enrichment of *Salmonella* in foodstuffs and other samples, according to ISO and FIL-IDF standards.

Presentation

20 Tubes
 Tube 16 x 113 mm
 with: 10 ± 0.3 ml

Packaging Details

16x113 mm glass tubes, ink labelled, metal-Non injectable cap. - 20 tubes per box

Shelf Life

10 months

Storage

8-25 °C

Composition

Composition (g/l):

Soy peptone.....	4.500
Sodium chloride.....	7.200
Monopotassium phosphate.....	1.260
Dipotassium phosphate.....	0.180
Magnesium chloride anhydrous.....	13.400
Malachite Green.....	0.036

Description /Technique

Description:

The Rappaport Vassiliadis medium complies with the recommendations of the APHA for the examination of food.

This culture medium is a modification of the R10 Medium (from Rappaport *et al.*) or RV Broth (from Vassiliadis *et al.*) by van Schothorst & Renaud. The modifications are an adjustment in the magnesium chloride concentration and the buffering capacity of the medium to aid pH maintenance during storage. It shows a higher selectivity towards *Salmonella* and produces better yields than other similar media, especially after preliminary enrichment and at an incubation temperature of 41 ± 0,5°C.

Malachite green, low pH and magnesium chloride inhibit the growth of microorganisms normally found in the intestine but do not affect the proliferation of most salmonellae. As malachite green inhibits the growth of *Shigella*, other culture methods may need to be used to isolate this organism. The addition of soy peptone enhances the growth of *Salmonella*.

Technique:

Collect, dilute and prepare samples and volumes as required according to specifications, directives, official standard regulations and/or expected results.

Inoculate aseptically the tubes with the prepared sample or its dilution in a proportion 1/10 (V/V).

Incubate the tubes tightly closed aerobically at 41 ± 0,5 ° C for 24h ± 3 h..

(Incubation times, temperature and sample volumes may vary depending on the sample, on the specifications,...)

Read the turbidity (growth indicator) and inoculate any confirmatory, secondary medium by streaking methodology or by spiral method, like XLD, BPLS,... to confirm results after proper incubation, enumerate all the colonies that have appeared onto the surface of the secondary agar.

Presumptive isolation / recovery of *Salmonella* must be confirmed by further microbiological and biochemical tests.

Each laboratory must evaluate the results according to their specifications.

Calculate total microbial count per ml of sample by multiplying the average number of colonies per plate by the inverse dilution factor if streaked a diluted sample. Report results as Colony Forming Unit (CFU's) per ml or g along with enrichment and secondary media used, incubation time and temperature.

Quality control

Physical/Chemical control

Color : Blue

pH: 5.2 ± 0.2 at 25°C

Microbiological control

Inoculate: Practical range 100 ± 20 CFU. min. 50 CFU (productivity)/ 10⁴-10⁶ (selectivity).

Microbiological control according to ISO 11133:2014/A1:2018.

Analytical methodology according to ISO 11133:2014/A1:2018; A2:2020

Aerobiosis. Incubation 41,5±1 °C, reading 24h ± 3h. Confirm in XLD (Pr) y TSA (Select.)

Microorganism

Enterococcus faecalis ATCC® 29212, WDCM 00087

Escherichia coli ATCC® 25922, WDCM 00013

S. typhimurium (14028) + *E. coli* (8739) + *Ps.* (27853)

Sterility Control

Incubation 48 h at 30-35 °C and 48 h at 20-25 °C: NO GROWTH.

Check at 7 days after incubation in same conditions.

Growth

Inhibited. Confirm in TSA at 37°C±1 reading 24 ± 3h

Partial Inhibition. Confirm in TSA at 37°C±1 reading 24 ± 3h.

Salmonella coln. charact. in XLD (37°C±1 / 24 ± 3h) ≥ 10 CFU.

Bibliography

- ATLAS, R.M., L.C. PARKS (1993) Handbook of Microbiological Media. CRC Press, Inc. London.
- DOWNES, F.P. & K. ITO (2001) Compendium of Methods for the Examination of Foods. 4th ed. APHA. Washington. USA.
- FDA (Food and Drug Administrations) US (1998) Bacteriological Analytical Manual. 8th ed. Revision A. AOAC Internacional. Gaithersburg. MD. USA.
- FIL-IDF 93:2001 Standard. Milk and Milk Products. Detection of *Salmonella*. Brussels.
- HORWITZ, W. (2000) Official Methods of Analysis of AOAC International. Gaithersburg. MD. USA.
- ISO Standard 6579-1 (2017) Microbiology of food chain - Horizontal method for the detection, enumeration and serotyping of *Salmonella* - Part 1 : Detection of *Salmonella* spp.
- ISO 6785:2001 Standard. Milk and Milk Products. Detection of *Salmonella*.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- RAPPAPORT, F., N. KONFORTI & B. NAVON (1956) A new enrichment medium for certain salmonellae. J. Clin. Pathol. 9:261-266.
- VAN SCHOTHORST, M. & A.M. RENAUD (1983) Dynamics of *Salmonella* isolation with modified Rappaport's Medium (R10). J. appl. Bact. 54:209-215.
- VASSILIADIS, P. (1983) The Rappaport Vassiliadis (RV) enrichment medium for the isolation of salmonellas: An overview. J. Appl. Bact. 54:69-76.
- VASSILIADIS, P., PATERAKI, EPAPAICONOMOU, N., PAPADAKIS, J.A.A., TICHPOULOS, D. (1976) Nouveau procédé d'enrichissement de *Salmonella*. Ann. Microbiol. (Inst. Pasteur) 127B (195-200).