

# Antibiotic Medium N° 23

Cat. 1386

Standard medium used for antibiotic testing.

## Practical information

Applications	Categories
Antibiotic Assay	General use

Industry: Pharmaceutical/Veterinary

## Principles and uses

Antibiotic Medium N° 23 is a standard medium used for the microbiological assay of antibiotics.

The activity (potency) of an antibiotic can be demonstrated under suitable conditions by its inhibitory effect on microorganisms. Reduction in antimicrobial activity may reveal changes not demonstrated by chemical methods. The antibiotic media are identified numerically with names assigned by Grove and Randall in "Assay methods of antibiotics". The use of standardized culture media and strict control of all test conditions are essential requirements in the microbiological assay of antibiotics in order to obtain satisfactory test results.

## Formula in g/L

Glucose	10	Bacteriological agar	13,5
Potassium dihydrogenphosphate	0,45	Yeast extract	2,5
Dipotassium hydrogen phosphate	0,69		

## Preparation

Suspend 27,14 grams of the medium in one liter of distilled water. Mix well and dissolve by heating with frequent agitation. Boil for one minute until complete dissolution. Sterilize in autoclave at 121 °C for 15 minutes.

## Instructions for use

- Liquefy the medium and inoculate it at a suitable temperature, for the example 48 °C to 50 °C, with a known quantity of a suspension of microorganism sensitive to the antibiotic to be examined.
- Agitate the mixture gently to produce a homogeneous distribution and immediately pour into Petry dishes a quantity of the inoculated medium to form a layer 2-5 mm thick. Alternatively, the medium may consist of 2 layers, only the upper layer being inoculated.
- Prepare a solution of the reference substance and of the antibiotic to be examined having known concentrations and presumed to be equal activity.
- Apply the solutions to the surface of the medium, for example, in sterile cylinders of porcelain, stainless steel, or in cavities prepared in the agar.
- The same volume of the solution must be added to each cylinder or cavity.
- Alternatively, use a sterile absorbent paper disc, impregnate the discs with the solutions of the reference substance or the solutions of the antibiotics to be examined and place on the surface of the agar.

## Quality control

Solubility	Appearance	Color of the dehydrated medium	Color of the prepared medium	Final pH (25°C)
w/o rests	Fine powder	Beige	Clear amber, slightly opalescent	5,6±0,1

## Microbiological test

Incubation conditions: (30±1 °C / 16-18 h).

Microorganisms	Specification
Bacillus subtilis ATCC 6633	Good growth

## Storage

---

Temp. Min.:2 °C  
Temp. Max.:25 °C

## Bibliography

---

Grove and Randall. Assay Methods of Antibiotics, Medical Encyclopedia Inc. New York 1955. United States Pharmacopoeia Convention. 1955. The United States Pharmacopoeia, 23rd Ed. Biological Tests and Assays, p. 1690-1696. The United States Pharmacopoeia Convention, Rockville, Md.  
European Pharmacopoeia 7.0