

## Specification

Highly nutrient liquid medium for general purpose use, formulated according to Pharmacopeial Harmonised Method.

## Presentation

2 Prepared Baas /5L  
Bags 5L  
with: 5000 ± 15 ml

### Packaging Details

1 box with 2 bags of 5L.  
PVC plasticizer free sterile bag with: 1 vial stopper + 1 penetrable cap.  
Dimensions: 27 x 40 cm. For use in food testing.

### Shelf Life

16 months

### Storage

2-25 °C

## Composition

Composition (g/l):

Peptone from casein .....	17.0
Soya peptone.....	3.00
Sodium chloride.....	5.00
Dipotassium phosphate.....	2.50
Dextrose.....	2.50

## Description /Technique

### Description :

Liquid culture medium used in Sterility Test, according to European pharmacopeia and USP.

Use according expected results, according type of samples and validated methods.

### Technique:

All the conditions and data for the validation of aseptic filling process can be consulted in the ISO Standard 13408-1 in the chapters devoted to the methods of preparation of sterile products in several Pharmacopoeias.

Each Bag is intended for use with an automatic dispenser in laboratories requiring large volumes of broth media or diluent.

Discard any partially used bag to avoid contamination.

The bag has multiple connection points: 1 penetrable cap (injection port) latex-free polycarbonate, for any additive injection required. And an injection (vial stopper) to connect to any standard equipment laboratory dosing with a connector.

Once completely empty, the bag can be disposed of along with normal plastic (PVC).

## Quality control

### Physical/Chemical control

Color : Yellowish-brown

pH: 7.3 ± 0.2 at 25°C

### Microbiological control

Prepare tubes - Inoculate: Practical range 10-100 CFU (productivity) according to harmonized Eur. Pharmacopoeia

Aerobiosis. Incubation at 30-35 °C. Read after 18-24 h to 72 h for bacteria and 3-5 days for fungi.

### Microorganism

*Bacillus subtilis* ATCC® 6633, WDCM 00003

*Ps. aeruginosa* ATCC® 9027, WDCM 00026

*Escherichia coli* ATCC® 8739, WDCM 00012

*Staphylococcus aureus* ATCC® 6538, WDCM 00032

*Salmonella typhimurium* ATCC® 14028, WDCM 00031

*Aspergillus brasiliensis* ATCC® 16404

*Candida albicans* ATCC® 10231, WDCM 00054

### Growth

Good

Good

Good

Good

Good

Good

Good

### Sterility Control

Incubation 14 days at 32.5 ± 2 °C: NO GROWTH.

Incubation 14 days at 22.5 ± 2 °C: NO GROWTH.

**Bibliography**

- ATLAS, R.M. & L.C. PARKS (1993) Handbook of Microbiological Media. CRC Press, Inc. London.
- DOWNES, F.P. & K. ITO (2001) Compendium of Methods for the Microbiological Examination of Food, 4th ed. ASM. Washington. DC.
- EUROPEAN PHARMACOPOEIA 8.0 (2014) 8th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- FDA (Food and Drug Administrations) (1998) Bacteriological Analytical Manual. 8th ed. Revision A. AOAC International. Gaithersburg. MD.
- HORWITZ, W. (2000) Official Methods of Analysis of AOAC INTERNATIONAL. 17th ed. Gaithersburg. MD. USA.
- ISO 11133:2014/ Adm 1:2018. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- ISO 9308-1 Standard (2000) Water Quality. Detection and enumeration of E. coli and coliform bacteria. Membrane filtration method.
- PASCUAL ANDERSON, M<sup>ª</sup>R<sup>ª</sup> (1992) Microbiología Alimentaria. Díaz de Santos S.A., Madrid.
- USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.
- USP 33 - NF 28 (2011) <71> Sterility Tests. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.