



Product: **SABOURAUD DEXTROSE AGAR DOUBLE WRAPPING IRRADIATED**

Specification

Solid medium for the enumeration and cultivation of fungi according to the Pharmacopeial Harmonised Method and ISO standard.

Presentation

	Packaging Details	Shelf Life	Storage
20 Plates /Irradiated 90 mm - Double wrapping with: 22 ± 1 ml	1 box with 2 cellophane bags (double wrapping) with 10 plates/bag. Every pack exhibitis a irradiation indicator stacked on the side of the bag.(8-14kGy).	3,5 months	2-14°C

Composition

Composition (g/l):

D(+)-Glucose.....	40.0
Peptone from casein	5.00
Meat Peptone.....	5.00
Agar.....	15.0

Description /Technique

Sabouraud Dextrose Agar is a modification of the classical Sabouraud medium for the cultivation of fungi. This new formula helps to maintain the morphology of fungi, providing a reliable medium for both cultivation and differentiation. Its selectivity is due to a low pH and a high glucose concentration, which together with incubation at a relatively lower temperature (20 -25°C) favours the growth of fungi while discouraging that of bacteria. The mixture of peptones employed has been selected to provide the fungi with all their nitrogen requirements.

Spread the plate streaking methodology or by spiral method.

Each laboratory must evaluate the results according to their specifications.

Attention: Petri plates are used for monitoring the microbiological contamination of surface and air inside cleanrooms, isolators, RABS, food industries and hospitals. The double/triple irradiated wrapping ensures that the package itself doesn't contaminate the environment as the first wrapper is removed just before entering the clean area.

Quality control

Physical/Chemical control

Color : Straw-coloured yellow pH: 5.6 ± 0.2 at 25°C

Microbiological control

Growth Promotion Test according to harmonized pharmacopoeial monographs and test methods & ISO 11133:2014

Spiral Spreading: Practical range 50 - 100 CFU (Productivity).

Aerobiosis. Incubation at 20-25°C. Reading ≤5 days.

Microorganism

Candida albicans ATCC® 10231, WDCM 00054

Aspergillus brasiliensis ATCC® 16404, WDCM 00053

S. cerevisiae ATCC® 9763, WDCM 00058

Growth

Good (≥70 %)

Good (≥70 %)

Good (≥70 %)

Sterility Control

Incubation 48 hours at 30-35°C and 48 hours at 20-25°C: NO GROWTH
Check at 7 days after incubation in same conditions



pronadisa
Micro & Molecular Biology

Reference: 0841

Technical Data Sheet

Product: **SABOURAUD DEXTROSE AGAR DOUBLE WRAPPING IRRADIATED**

Bibliography

- AJELLO, L. (1957) Cultural Methods for Human Pathogenic Fungi J. Chron. Dis. 5:545-551.
- COLIPA (1997) Guidelines on Microbial Quality Management (MQM). Brussels.
- EUROPEAN PHARMACOPOEIA 8.0 (2014) 8th ed. § 2.6.13. Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. EDQM. Council of Europe. Strasbourg.
- GEORGE, L.K., AJELLO, L. & PAPAGEORGE, C. (1954) Use of Cycloheximide in the Selective Isolation of Fungi Pathogenic to Man. J. Lab. Clin. Med, 44 (422-428).
- HANTSCHKE, D. (1968) Mykosen, 11, (769-778).
- ISO 11133:2014. Microbiology of food, animal feed and water. Preparation, production, storage and performance testing of culture media.
- PAGANO, J. LEVIN, J.D. and TREJO, W. (1957-58) Diagnostic Medium for Differentiation of Species of *Candida*. Antibiotics Annual, 137-143.
- SABOURAUD, R. (1910) Les Teignes. Masson, Paris.
- USP 33 - NF 28 (2011) <62> Microbiological examination of non-sterile products: Test for specified microorganisms. Harmonised Method. USP Corp. Inc. Rockville. MD. USA.